

R_b and New Physics: A Comprehensive Analysis

J.M. CLINE

McGill University, 3600 University St., Montréal, Québec H3A 2T8, Canada

I summarize work¹ done with Bamert, Burgess, Nardi and London in which we examine the effects of new physics on R_b . Our work facilitates the identification of such new physics in case the present two-sigma deviation of R_b from its Standard Model value should persist, or alternatively it helps to constrain new physics in case the discrepancy continues to disappear. We consider two general classes of effects: mixing of the b or t quarks with heavy, exotic quarks, and loop corrections to the $Zb\bar{b}$ vertex. Our aim was to give an analysis which is as exhaustive and as model-independent as possible.

1 The (late?) R_b problem

As of one day before this talk was given, there appeared to be a serious discrepancy with the LEP measurement of R_b , the ratio of the partial widths $\Gamma(Z \rightarrow b\bar{b})$ versus $\Gamma(Z \rightarrow \text{hadrons})$. Naturally, this inspired a plethora of possible theoretical explanations. In trying to judge the relative merits of one extension of the Standard Model versus another, it would be useful to have some kind of unified framework in which to view them, so as to be able to answer questions like “how special is this particular model?” or “could another similar model do the job better?” We were thus led to consider the more general question “What kind of new physics can increase the $Zb\bar{b}$ vertex?” and to try to answer it in as model-independent a way as possible.

Let me first review the experimental situation as it stood just after the Moriond 1996 conference.² Normalizing the $Zb\bar{b}$ couplings as

$$\frac{e}{\sin 2\theta_w} \bar{b} \gamma^\mu (g_L^b(1 - \gamma_5) + g_R^b(1 + \gamma_5)) b Z_\mu, \quad (1)$$

their Standard Model values can be expressed as

$$\begin{aligned} g_L^b &= -\frac{1}{2} + \frac{1}{3} \sin^2 \theta_w = -0.4230; \\ g_R^b &= +\frac{1}{3} \sin^2 \theta_w = +0.770. \end{aligned} \quad (2)$$

Allowing for deviations δg_L^b and δg_R^b of these couplings from their Standard Model values, possibly due to new physics, and performing a global fit to the Z -pole data, one obtains the results³ summarized in table 1. Column 1 shows which parameters among δg_L^b and δg_R^b are allowed to vary, so that row 1 (where neither vary) represents the Standard Model, which is a poor fit to the data.

Table 1: Global fits to Moriond Z -pole data.³

vary	δg_L^b	δg_R^b	c.l. of fit	$\chi^2/\text{d.o.f}$
none	0	0	2.5%	24.7/13
δg_L^b	-0.0063 ± 0.0020	0	26%	14.7/12
δg_R^b	0	0.034 ± 0.010	37%	13.0/12
both	-0.0029 ± 0.0037	0.022 ± 0.018	32%	12.5/11

This is due to the measured $R_b = 0.2219 \pm 0.0017$ having been larger than the Standard Model value of $R_b^{SM} = 0.2156$. Allowing δg_L^b or δg_R^b to vary increases the confidence level of the fit from 2.5% to 26 – 37%, a quite substantial improvement. From a statistical point of view, it is sufficient to adjust either of the couplings, not necessarily both, to get an acceptable fit.

Table 1 shows that it takes a much bigger relative change in the right-handed than the left-handed coupling to change R_b by the amount that was needed: $\delta g_R^b/g_R^b \sim 40\%$ versus $\delta g_L^b/g_L^b \sim 2\%$. This immediately tells us that we need tree-level physics, for example mixing of b_R with new quarks, to get a big enough change in g_R^b . On the other hand, one-loop effects or small tree-level mixing can give a sufficiently large change in the left-handed coupling. There is a further possibility we do not discuss, but which has been pursued by others,⁴ mixing of the Z boson with a Z' .

2 Quark mixing*2.1 b - b' mixing*

A simple way to change the $Zb\bar{b}$ couplings is to allow mixing between the b and a new exotic b' with both left- and right-handed components.

Table 2: Twelve models of b - b' mixing that solve the R_b problem, labeled by the isospins of the b'_L and b'_R .

I'_L	I'_{3L}	I'_R	I'_{3R}	angle needed
1	-1	1	-1	$s_L^2 = 0.0111 \pm 0.0032$
1	-1	1/2	-1/2	"
3/2	-3/2	1	-1	$s_L^2 = 0.0056 \pm 0.0016$
1/2	1/2	1/2	1/2	$s_R^2 = 0.052^{+0.013}_{-0.014}$
0	0	1/2	1/2	"
1/2	1/2	1	1	$s_R^2 = 0.026^{+0.006}_{-0.007}$
1/2	1/2	0, 1	0	$s_L^2 = 0.8515 \pm 0.0016$
3/2	1/2	1	0	"
0	0	1/2	-1/2	$s_R^2 \gtrsim 0.361$
1/2	-1/2	1/2	-1/2	$s_R^2 = 0.361^{+0.013}_{-0.014}$
1/2	-1/2	1	-1	$s_R^2 = 0.180 \pm 0.007$

The couplings become $g_{L,R}^b \rightarrow \cos^2 \theta_{L,R} g_{L,R}^b + \sin^2 \theta_{L,R} g_{L,R}^{b'}$, where the mixing angles come from the similarity transformation $M \rightarrow O_L^T M O_R$ needed to diagonalize the 2×2 Dirac mass matrix M for b and b' . We were able to simplify the range of possibilities by making three reasonable assumptions. First of all one must assume that M_{12} or M_{21} is nonvanishing in order to have any mixing at all. Second, we expect the b' to be heavy, which is only possible if M_{22} , the entry corresponding to the b' mass in the case of zero mixing, is large. Finally we exclude Higgs boson representations higher than doublets, since these tend to make undesirably large contributions to the rho parameter.

Under these assumptions there are only twelve possible isospin assignments for the chiral components of the b' quark, which are listed in Table 2, along with the mixing angle needed ($s_{L,R}^2 \equiv \sin^2 \theta_{L,R}$) to change g_L^b or g_R^b by the desired amount. In almost all of these models, only one of the two mixing angles is relevant, the other one being suppressed by powers of the first. However for the model with $I'_L = I'_{3L} = 0$ and $I'_R = -I'_{3R} = 1/2$, a combination of both left- and right-handed mixing angles is possible, illustrated in figure 1.

2.2 t - t' mixing

Another possibility is that the top quark mixes with an exotic t' , which alters the g_L^b coupling




Figure 1: Allowed values of mixing angles in the s_R^2 - s_L^2 plane for model number 10 of table 2.




Figure 2: Top quark loop diagrams that affect g_L^b .

through the loop diagrams of figure 2. These are the same as the diagrams of the Standard Model, except that the top quark must be replaced by two linear combinations of t and t' , one for each chirality. Making the same assumptions about the t - t' mass matrix as we did for that of b and b' above, there are again twelve possible isospin assignments for the t' quark, enumerated in Table 3.

The resulting change in g_L^b can be expressed rather simply in the limit where both top quarks are much heavier than the W boson, if we concentrate on values of $m_{t'}$ so as to maximize the magnitude of g_L^b . This turns out to occur when $m_{t'} \ll m_t$, in which case a single term dominates

Table 3: Twelve possible models of t - t' mixing, labeled by the isospins of the t'_L and t'_R .

I'_L	I'_{3L}	I'_R	I'_{3R}
3/2	+1/2	1	+1
1/2	+1/2	1	+1, 0
1/2	+1/2	1/2	+1/2
1/2	+1/2	0	0
1/2	-1/2	1	0, -1
1/2	-1/2	1/2	-1/2
1/2	-1/2	0	0
0	0	1/2	$\pm 1/2$
0	0	0	0

the shift in g_L^b beyond its Standard Model value,

$$\delta g_L^b \cong \frac{\alpha}{16\pi s_w^2} V_{t'b}^2 \left(\frac{m_{t'}^2 - m_t^2}{m_W^2} + 6 \ln \frac{m_{t'}}{m_t} \right). \quad (3)$$

Here $V_{t'b}$ is the element from the extended CKM matrix which includes the t' quark in addition to u , c and t . The best case is when $m_{t'}$ is at its experimental lower limit of 135 GeV, in which case $\delta g_L^b = -0.0021$. Although this is too small to explain the previous LEP data which needed $\delta g_L^b = -0.0063$, the recent results of ALEPH announced at this conference⁵ have reduced the size of the discrepancy between theory and the world average measurement⁶ of $R_b = 0.2178 \pm 0.0011$ such that -0.0021 is now sufficient.

3 General Loop Effects

3.1 Diagonal Z Couplings

In the previous subsection we had to consider loops involving the t' as a special case because they were inextricably tangled up with usual Standard Model contributions, but here we want to consider the effects of a general fermion f and scalar ϕ coupling to the b_L via the interaction $\mathcal{L}_y = y \bar{b}_L \phi f + \text{h.c.}$, and to the Z boson via

$$\frac{2e}{\sin 2\theta_W} \left(\bar{f} \gamma_\mu (g_L^f P_L + g_R^f P_R) f + i g^\phi \phi^\dagger \overleftrightarrow{\partial}_\mu \phi \right) Z^\mu. \quad (4)$$

The relevant diagrams which modify g_L^b are shown in figure 3. There are also vacuum polarization diagrams as in figure 4, but these cancel to a good approximation in the ratio that defines R_b . The result of evaluating the first four diagrams has a




Figure 3: The important generic loops for g_L^b




Figure 4: Vacuum polarization contribution which tend to cancel in R_b .

remarkably simple expression in the approximation of ignoring M_Z :

$$\delta g_L^b = \frac{y^2 n_c}{16\pi^2} (g_L^f - g_R^f) \mathcal{F}(m_f^2/m_\phi^2), \quad (5)$$

where n_c is a color factor that is unity if ϕ is colorless and f is a color triplet, for example. The kinematic function given by $\mathcal{F}(x) = x/(x-1) - x \ln x/(x-1)^2$ is always positive and reaches its maximum value of 1 when $m_f/m_\phi \rightarrow \infty$. The $M_Z = 0$ approximation turns out to be better than one would expect, since the first order correction in powers of M_Z^2/m_f^2 is typically only 10% of eq. (5), even when $M_Z^2/m_f^2 = 1$.

A notable feature of eq. (5) is that its sign is completely determined by the isospins of f_L and f_R since $g_L^f - g_R^f = I_{3L}^f - I_{3R}^f$. Thus one can immediately see why two-Higgs doublet models, as well as the top quark contribution in the Standard Model, tend to decrease R_b : if f is the top quark then $I_{3L}^f - I_{3R}^f = 1/2$, which has the opposite sign to the tree level value of g_L^b , reducing the effective coupling. In two-Higgs models with very large $\tan \beta$, so that the bottom quark itself can make an appreciable contribution in the loop, the sign is such as to increase R_b .

3.2 Nondiagonal Z Couplings

It might happen that the flavor states with the largest coupling to b_L , which we shall now denote by ϕ_1 and f_1 , mix with other flavor states ϕ_2 and f_2 to form mass eigenstates ϕ, ϕ' and f, f' . Then our previous formula (5) for δg_L^b must be generalized to something more complicated. Nevertheless

in certain limiting cases it becomes relatively simple again. First, if the masses are degenerate such that $m_f = m_{f'}$ and $m_\phi = m_{\phi'}$, we get

$$\delta g_L^b = \frac{y^2 n_c}{16\pi^2} (U_R U_L^\dagger g_L^f U_L U_R^\dagger - g_R^f)_{11} \mathcal{F}(m_f^2/m_\phi^2). \quad (6)$$

Here $g_{L,R}^f$ are the 2×2 matrix generalizations of the fermion couplings to the Z boson, and $U_L^\dagger M U_R$ is the similarity transformation that diagonalizes the f_1 - f_2 mass matrix. Second, if the fermions are much heavier than the scalars, we get eq. (6) again, with the addition of two extra terms that tend to cancel each other, at least in the supersymmetric case we will discuss below. Third, if the scalars are much heavier than the fermions we obtain

$$\delta g_L^b = \frac{y^2 n_c}{16\pi^2} \sin^2 2\theta_\phi (g_{11}^\phi - g_{22}^\phi) \mathcal{F}'(m_f^2/m_\phi^2), \quad (7)$$

where θ_ϕ is the scalar mixing angle and $\mathcal{F}'(x) = \frac{x+1}{2(x-1)} \ln x - 1$ is another positive function like $\mathcal{F}$. Thus it is once again rather easy to determine the sign and magnitude of δR_b .

An example is the supersymmetric Standard Model where, because of the large top quark yukawa coupling, ϕ_1 is the right-handed top squark $\tilde{t}_R$, and f_1 is the higgsino $\tilde{h}_2^-$, or alternatively the fermion and scalar are the usual top quark and one of the Higgs bosons, $\phi_1 = h_2^-$ and $f_1 = t_R$. We already explained that the quark-Higgs contribution has the wrong sign for increasing R_b , so one wants to minimize these loops by taking the Higgs bosons to be very heavy. Concentrating on the squark and higgsino, therefore, one must take into account their mixing with the left-handed top squark, $\phi_2 = \tilde{t}_L$ and the Wino, $f_2 = \tilde{W}^-$. The charge matrices are $g_L^f = g_R^f = \text{diag}(-1/2, -1)$ for the fermions and $g^\phi = \text{diag}(0, 1/2)$ for the scalars. Applying the simplifying cases 1 or 2 mentioned above, one finds

$$\delta g_L^b \cong -\frac{y^2 n_c}{32\pi^2} \sin^2(\theta_L - \text{sign}(m_f/m_{f'})\theta_R) \mathcal{F}, \quad (8)$$

and therefore to have a large shift in g_L^b , the combination of chargino mixing angles $|\theta_L - \text{sign}(m_f/m_{f'})\theta_R|$ must be large. From this and the form of the chargino mass matrix it is easy to deduce that both charginos must be rather light, and that $\tan\beta$ must be close to unity. Furthermore the third simplifying limit above, that of

heavy scalars, gives the wrong sign for increasing R_b because $g_{11}^\phi - g_{22}^\phi = -1/2$. Thus one needs at least one light scalar, which naturally enough turns out to be $\tilde{t}_R$ because this is the one that couples directly to the left-handed b quark.

4 Summary

We have shown that a significant increase of R_b from its Standard Model value can be caused by new physics that changes g_R^b by quark mixing, or g_L^b either by mixing or one loop contributions to the $Zb\bar{b}$ coupling. In the category of b - b' mixing we found 12 possible models using modest restrictions. The Standard Model t quark loop corrections to g_L^b can also be altered by t - t' mixing, but only by as much as -0.0021 , corresponding to a maximum increase of 0.0017 in R_b .

For more general kinds of loop corrections to R_b , we have found simple approximate formulas which gives insight into the sign and magnitude of the change. Although the original motivation was to make it easier to build models explaining the former discrepancy, perhaps they will in the future be more useful in constraining models that arise in other contexts. It should however be kept in mind that the world average value of R_b as of this writing is still two standard deviations higher than the Standard Model value, despite the recent results of ALEPH which now agree with the Standard Model.

References

1. P. Bamert, C.P. Burgess, J.M. Cline, E. Nardi and D. London, *Phys. Rev. D* **54**, 4275 (1996).
2. LEP electroweak working group, preprint CERN-PPE/95-172, 1995; M.D. Hildreth, "The Current Status of the R_b and R_c Puzzle," talk given at the XXXIst Rencontres de Moriond, March 1996.
3. P. Bamert, preprint McGill-95-64, hep-ph/9512445 (1996).
4. see for example M. Mangano, "Hadrophilic/Leptophobic Z' Models and R_b ," in these proceedings.
5. I. Tomalin, " R_b with Multivariate Analysis," in these proceedings.
6. K. Mönig, "Combination of Heavy Quark E.W. measurements," in these proceedings.